

58. (New) A method for breaking B-cell autotolerance in an animal to a self-protein of that animal, and inducing antibody production in the animal against the self-protein of that animal, comprising administering to the animal, an immunologically effective amount of at least one modified self-protein, wherein:

the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving tertiary structure of the self-protein, and said substitution preserving flanking regions comprising at least four amino acids on each side of the peptide fragment;

whereby the modified self-protein elicits antibodies that are against the self-protein, and B-cell autotolerance to the self-protein is broken.

59. (New) A method for breaking B-cell autotolerance in an animal to a self-protein of that animal, and inducing antibody production in the animal against the self-protein of that animal, comprising administering to the animal, an immunologically effective amount of at least one modified self-protein, wherein:

the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving tertiary structure of the self-protein, and said substitution preserving flanking regions comprising at least ten amino acids on each side of the peptide fragment;

whereby the modified self-protein elicits antibodies that are against the self-protein, and B-cell autotolerance to the self-protein is broken.

60. (New) A method for breaking B-cell autotolerance in an animal to a self-protein of that animal, and inducing antibody production in the animal against the self-protein of that animal, comprising administering to the animal, an immunologically effective amount of at least one modified self-protein, wherein:

the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving tertiary structure of the self-protein, and said substitution preserving flanking regions comprising at least fifteen amino acids on each side of the peptide fragment,

whereby the modified self-protein elicits antibodies that are against the self-protein, and B-cell autotolerance to the self-protein is broken.

61. (New) A method for breaking B-cell autotolerance in an animal to a self-protein of that animal, and inducing antibody production in the animal against the self-protein of that animal, comprising administering to the animal, an immunologically effective amount of at least one modified self-protein, wherein:

the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving tertiary structure of the self-protein,

whereby the modified self-protein elicits antibodies that are against the self-protein; and, the modified self-protein elicits an immune response in the animal which includes an MHC class II immune response as to the immunodominant T-cell epitope and an autoantibody response in other MHC-haplotypes, and B-cell autotolerance to the self-protein is broken.

62. (New) A method for breaking B-cell autotolerance in an animal to a self-protein of that animal, and inducing antibody production in the animal against the self-protein of that animal, comprising administering to the animal, an immunologically effective amount of at least one modified self-protein, wherein:

the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by being detoxified and by containing a substitution of at least one peptide fragment of the self-protein

with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving tertiary structure of the self-protein, whereby the modified self-protein elicits antibodies that are against the self-protein, and B-cell autotolerance to the self-protein is broken.

63. (New) A method for breaking B-cell autotolerance in an animal to a self-protein of that animal, and inducing antibody production in the animal against the self-protein of that animal, comprising administering to the animal, an immunologically effective amount of at least one modified self-protein, wherein:

the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving secondary and tertiary structure of the self-protein, and said substitution preserving flanking regions comprising at least four amino acids on each side of the peptide fragment;

whereby the modified self-protein elicits antibodies that are against the self-protein, and B-cell autotolerance to the self-protein is broken.

64. (New) A method for breaking B-cell autotolerance in an animal to a self-protein of that animal, and inducing antibody production in the animal against the self-protein of that animal, comprising administering to the animal, an immunologically effective amount of at least one modified self-protein, wherein:

the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving secondary and tertiary structure of the self-protein, and said substitution preserving flanking regions comprising at least ten amino acids on each side of the peptide fragment;

whereby the modified self-protein elicits antibodies that are against the self-protein, and B-cell autotolerance to the self-protein is broken.

65. (New) A method for breaking B-cell autotolerance in an animal to a self-protein of that animal, and inducing antibody production in the animal against the self-protein of that animal, comprising administering to the animal, an immunologically effective amount of at least one modified self-protein, wherein:

the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving secondary and tertiary structure of the self-protein, and said substitution preserving flanking regions comprising at least fifteen amino acids on each side of the peptide fragment,

whereby the modified self-protein elicits antibodies that are against the self-protein, and B-cell autotolerance to the self-protein is broken.

66. (New) A method for breaking B-cell autotolerance in an animal to a self-protein of that animal, and inducing antibody production in the animal against the self-protein of that animal, comprising administering to the animal, an immunologically effective amount of at least one modified self-protein, wherein:

the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving secondary and tertiary structure of the self-protein,

whereby the modified self-protein elicits antibodies that are against the self-protein; and, the modified self-protein elicits an immune response in the animal which includes an MHC class II immune response as to the immunodominant T-cell epitope and an autoantibody response in other MHC-haplotypes, and B-cell autotolerance to the self-protein is broken.

67. (New) A method for breaking B-cell autotolerance in an animal to a self-protein of that animal, and inducing antibody production in the animal against the self-protein of that animal, comprising administering to the animal, an immunologically effective amount of at least one modified self-protein, wherein:

the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by being detoxified and by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving secondary and tertiary structure of the self-protein,

whereby the modified self-protein elicits antibodies that are against the self-protein, and B-cell autotolerance to the self-protein is broken.

68. (New) A method for breaking B-cell autotolerance in an animal to a self-protein of that animal, and inducing antibody production in the animal against the self-protein of that animal, comprising;

preparing different modified self-proteins, wherein:

each modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal,

said substitution preserving tertiary structure of the self-protein, and

the different modified self-proteins differ from each other with respect to the position of the at least one immunodominant T-cell epitope;

ascertaining which of the different modified self-proteins elicits a desired specific neutralizing effect and thereby ascertaining a desired modified self-protein; and

administering to the animal, an immunologically effective amount of the desired modified self-protein, wherein:

the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein, and,

the desired modified self-protein elicits antibodies that are against the self-protein, and B-cell autotolerance to the self-protein is broken.

69. (New) A method for breaking B-cell autotolerance in an animal to a self-protein of that animal, and inducing antibody production in the animal against the self-protein of that animal, comprising;

preparing different modified self-proteins, wherein:

each modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving secondary and tertiary structure of the self-protein, and

the different modified self-proteins differ from each other with respect to the position of the at least one immunodominant T-cell epitope;

ascertaining which of the different modified self-proteins elicits a desired specific neutralizing effect and thereby ascertaining a desired modified self-protein; and

administering to the animal, an immunologically effective amount of the desired modified self-protein, wherein:

the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein, and,

the desired modified self-protein elicits antibodies that are against the self-protein, and B-cell autotolerance to the self-protein is broken.

70. (New) A method for breaking B-cell autotolerance in an animal to a self-protein of that animal, and inducing antibody production in the animal against the self-protein of that animal, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, comprising administering to the animal, an immunologically effective amount of at least one modified self-protein, wherein:

a. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving tertiary structure of the self-protein;

whereby, the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

b. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving secondary and tertiary structure of the self-protein;

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

c. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving tertiary structure of the self-protein, and said substitution preserving flanking regions comprising at least four amino acids on each side of the peptide fragment;

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

d. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving tertiary structure of the self-protein, and said substitution preserving flanking regions comprising at least ten amino acids on each side of the peptide fragment;

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier

protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

e. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving tertiary structure of the self-protein, and said substitution preserving flanking regions comprising at least fifteen amino acids on each side of the peptide fragment,

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

f. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving tertiary structure of the self-protein,

whereby the modified self-protein elicits antibodies that are against the self-protein earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes; and, the modified self-protein elicits an immune response in the animal which includes an MHC class II immune response as to the immunodominant T-cell epitope and an autoantibody response in other MHC-haplotypes, and B-cell autotolerance to the self-protein is broken; or,

g. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by being detoxified and by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving tertiary structure of the self-protein,



whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

h. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving secondary and tertiary structure of the self-protein, and said substitution preserving flanking regions comprising at least four amino acids on each side of the peptide fragment;

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

i. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving secondary and tertiary structure of the self-protein, and said substitution preserving flanking regions comprising at least ten amino acids on each side of the peptide fragment;

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

j. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self- protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving secondary and tertiary structure of the self-protein, and said substitution preserving flanking regions comprising at least fifteen amino acids on each side of the peptide fragment,

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

k. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self- protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving secondary and tertiary structure of the self-protein,

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes; and, the modified self-protein elicits an immune response in the animal which includes an MHC class II immune response as to the immunodominant T-cell epitope and an autoantibody response in other MHC-haplotypes, and B-cell autotolerance to the self-protein is broken; or,

1. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self- protein, by being detoxified and by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving secondary and tertiary structure of the self-protein,

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier

protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

m. preparing different modified self-proteins, wherein:

each modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal,

said substitution preserving tertiary structure of the self-protein, and the different modified self-proteins differ from each other with respect to the position of the at least one immunodominant T-cell epitope;

ascertaining which of the different modified self-proteins elicits a desired specific neutralizing effect and thereby ascertaining a desired modified self-protein; and

administering to the animal, an immunologically effective amount of the desired modified self-protein, wherein:

the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein, and,

the desired modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

n. preparing different modified self-proteins, wherein:

each modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal,

said substitution preserving secondary and tertiary structure of the self-protein, and

the different modified self-proteins differ from each other with respect to the position of the at least one immunodominant T-cell epitope;

ascertaining which of the different modified self-proteins elicits a desired specific neutralizing effect and thereby ascertaining a desired modified self-protein; and

administering to the animal, an immunologically effective amount of the desired modified self-protein, wherein:

the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein, and,

the desired modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken.

71. (New) A method for breaking B-cell autotolerance in an animal to a self-protein of that animal, inducing antibody production in the animal against the self-protein of that animal, and eliciting an immune response in the animal which includes an MHC class II immune response as to an immunodominant T-cell epitope which is foreign to the animal and an autoantibody response in other MHC-haplotypes, comprising administering to the animal, an immunologically effective amount of at least one modified self-protein, wherein:

a. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing the immunodominant T-cell epitope which is foreign to the animal, said substitution preserving tertiary structure of the self-protein,

whereby the modified self-protein elicits antibodies that are against the self-protein; and, the modified self-protein elicits an immune response in the animal which includes an MHC class II immune response as to the immunodominant T-cell epitope and an autoantibody response in other MHC-haplotypes, and B-cell autotolerance to the self-protein is broken; or

b. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing the immunodominant T-cell epitope which is foreign to the animal, said substitution preserving secondary and tertiary structure of the self-protein,

whereby the modified self-protein elicits antibodies that are against the self-protein; and, the modified self-protein elicits an immune response in the animal which includes

K1 Cont  
an MHC class II immune response as to the immunodominant T-cell epitope and an autoantibody response in other MHC-haplotypes, and B-cell autotolerance to the self-protein is broken.

72. (New) A method for breaking B-cell autotolerance in an animal to a self-protein of that animal, and inducing antibody production in the animal against the self-protein of that animal, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, comprising administering to the animal, an immunologically effective amount of at least one modified self-protein, wherein:

a. the self-protein is normally non-immunogenic in the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving tertiary structure of the self-protein;

whereby, the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

b. the self-protein is normally non-immunogenic in the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving secondary and tertiary structure of the self-protein;

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

c. the self-protein is normally non-immunogenic in the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide

containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving tertiary structure of the self-protein, and said substitution preserving flanking regions comprising at least four amino acids on each side of the peptide fragment;

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

d. the self-protein is normally non-immunogenic in the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving tertiary structure of the self-protein, and said substitution preserving flanking regions comprising at least ten amino acids on each side of the peptide fragment;

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

e. the self-protein is normally non-immunogenic in the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving tertiary structure of the self-protein, and said substitution preserving flanking regions comprising at least fifteen amino acids on each side of the peptide fragment,

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

f. the self-protein is normally non-immunogenic in the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self- protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving tertiary structure of the self-protein,

whereby the modified self-protein elicits antibodies that are against the self-protein earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes; and, the modified self-protein elicits an immune response in the animal which includes an MHC class II immune response as to the immunodominant T-cell epitope and an autoantibody response in other MHC-haplotypes, and B-cell autotolerance to the self-protein is broken; or,

g. the self-protein is normally non-immunogenic in the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self- protein, by being detoxified and by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving tertiary structure of the self-protein,

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

h. the self-protein is normally non-immunogenic in the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self- protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving secondary and tertiary structure of the self-protein, and said substitution preserving flanking regions comprising at least four amino acids on each side of the peptide fragment;

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier

protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

i. the self-protein is normally non-immunogenic in the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving secondary and tertiary structure of the self-protein, and said substitution preserving flanking regions comprising at least ten amino acids on each side of the peptide fragment;

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

j. the self-protein is normally non-immunogenic in the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving secondary and tertiary structure of the self-protein, and said substitution preserving flanking regions comprising at least fifteen amino acids on each side of the peptide fragment,

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

k. the self-protein is normally non-immunogenic in the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide



containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving secondary and tertiary structure of the self-protein,

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes; and, the modified self-protein elicits an immune response in the animal which includes an MHC class II immune response as to the immunodominant T-cell epitope and an autoantibody response in other MHC-haplotypes, and B-cell autotolerance to the self-protein is broken; or,

1. the self-protein is normally non-immunogenic in the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by being detoxified and by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving secondary and tertiary structure of the self-protein,

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

m. preparing different modified self-proteins, wherein:

each modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal,

said substitution preserving tertiary structure of the self-protein, and

the different modified self-proteins differ from each other with respect to the position of the at least one immunodominant T-cell epitope;

ascertaining which of the different modified self-proteins elicits a desired specific neutralizing effect and thereby ascertaining a desired modified self-protein; and

administering to the animal, an immunologically effective amount of the desired modified self-protein, wherein:

the self-protein is normally non-immunogenic in the animal and there is normally B-cell autotolerance by the animal to the self-protein, and,

the desired modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

n. preparing different modified self-proteins, wherein:

each modified self-protein is modified, in comparison to the self-protein, by containing a substitution of at least one peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, said substitution preserving secondary and tertiary structure of the self-protein, and

the different modified self-proteins differ from each other with respect to the position of the at least one immunodominant T-cell epitope;

ascertaining which of the different modified self-proteins elicits a desired specific neutralizing effect and thereby ascertaining a desired modified self protein; and

administering to the animal, an immunologically effective amount of the desired modified self-protein, wherein:

the self-protein is normally non-immunogenic in the animal and there is normally B-cell autotolerance by the animal to the self-protein, and,

the desired modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken.

73. (New) The method of any one of claims 56-72 wherein the modified self-protein is a recombinant modified self-protein.

74. (New) The method of any one of claims 56-72 wherein the self-protein is tumor necrosis factor alpha (TNF- $\alpha$ ), tumor necrosis factor beta (TNF- $\beta$ ), gamma interferon ( $\gamma$ -interferon), interleukin 1 (IL-1) or immune globulin (IgE).

75. (New) The method of claim 73 wherein the self-protein is tumor necrosis factor alpha (TNF- $\alpha$ ), tumor necrosis factor beta (TNF- $\beta$ ), gamma interferon ( $\gamma$ -interferon), interleukin 1 (IL-1) or immune globulin (IgE).

Sub 1 K3  
76. (New) The method of any one of claims 56-72 wherein the administering includes administering an adjuvant.

77. (New) The method of claim 76 wherein the adjuvant comprises calcium phosphate, saponin, quil A or a biodegradable polymer.

78. (New) The method of claim 73 wherein the administering includes an adjuvant.

79. (New) The method of claim 75 wherein the administering includes an adjuvant.--

---

Kindly cancel claims 26, 28-43, 45-48 and 51-55, without prejudice, without admission, without surrender of subject matter, and without any intention of creating any estoppel as to equivalents.